

### Outlineoffungi.org - Note 3

*Amerosporiopsis*, *Amphiportha*, *Anaphysmene*, *Anthracoderma*, *Alloneottiosporina*, *Aoria*, *Aphanofalx*, *Apiocarpella* and *Apiognomonina*

Li et al. (2020) treated all eight genera and provided descriptions and drawings. No new collections came to light and *Amerosporiopsis*, *Anaphysmene*, *Anthracoderma*, *Alloneottiosporina*, *Aoria*, *Aphanofalx* and *Apiocarpella* remain in Ascomycota genera *incertae sedis* pending fresh collections. *Amphiportha* Petr. and *Apiognomonina* Höhn. Are confirmed as genera of Gnomoniaceae in Diaporthales, Sordariomycetes.

#### Reference

Li WJ, McKenzie EHC, Liu JK, Bhat J, Dai DQ, Camporesi E, Tian Q, Maharachchikumbura SSN, Luo ZL, Shang QJ, Zhang FJ, Tangthirasunun N, Karunarathna SC, Xu JC, Hyde KD (2020) Taxonomy and phylogeny of hyaline-spored coelomycetes. *Fungal Diversity* 100, 279–801.

#### Entry by

**Kevin D. Hyde**, Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai, Thailand

(Edited by: **Nalin Wijayawardene**)