## Outlineoffungi.org - Note 995 Barbatosphaeriales

Web-links: Index Fungorum, Facesoffungi, MycoBank, GenBank

## **Barbatosphaeriales** K.D. Hyde & Hongsanan

<u>Barbatosphaeriaceae</u> was introduced by Zhang et al. (2017) to accommodate three genera, <u>Barbatosphaeria</u>, <u>Ceratostomella</u> and <u>Xylomelasma</u>, as they share similar characteristics, such as dark, long-necked, astromatic ascomata, clavate or cylindrical-clavate asci and mostly ellipsoidal ascospores. In addition, Conidiophores are cylindrical to fask- or irregularly-shaped, and brown or hyaline. Conidiogenous cells are polyblastic, and integrated, terminal. Conidia are unicellular and hyaline. Hyde et al. (2020) placed <u>Barbatosphaeriaceae</u> in <u>Diaporthomycetidae</u> families <u>incertae sedis</u> based on phylogenetic analysis. Hyde et al. (2021) raised <u>Barbatosphaeriaceae</u> to order status based on phylogenetic analysis coupled with divergent time estimation. <u>Barbatosphaeria</u> Réblová, as the type genus, is a non-stromatic perithecial ascomycete introduced for a single species <u>Barbatosphaeria barbirostris</u> (Dufour) Réblová (<u>Réblová 2007</u>). The members of this order are saprobic on decaying wood or plant material. The taxonomic placement of <u>Barbatosphaeriales</u> is in <u>Diaporthomycetidae</u>, <u>Sordariomycetes</u>, <u>Pezizomycotina</u>, and <u>Ascomycota</u>.

## References

- Réblová 2007 *Barbatosphaeria* gen. et comb. nov., a new genus for *Calosphaeria barbirostris*. Mycologia 99, 723–732. https://doi.org/10.1080/15572536.2007.11832536
- Zhang H, Dong W, Hyde KD, Maharachchikumbura SSN, Hongsanan S et al. 2017 Towards a natural classification of Annulatascaceae-like taxa: introducing Atractosporales ord. nov. and six new families. Fungal Diversity 85, 75–110. https://doi.org/10.1007/s13225-017-0387-z
- Hyde KD, Norphanphoun C, Maharachchikumbura SSN, Bhat DJ, Jones EBG et al. 2020 Refned families of *Sordariomycetes*. Mycosphere 11, 305–1059. https://doi.org/10.5943/mycosphere/11/1/7
- Hyde KD, Bao DF, Hongsanan S, Chethana KWT et al. 2021 Evolution of freshwater *Diaporthomycetidae* (*Sordariomycetes*) provides evidence for five new orders and six new families. Fungal Diversity 107, 71–105. https://doi.org/10.5943/mycosphere/11/1/7

## **Entry by**

**Zong-Long Luo**, College of Agriculture and Biological Science, Dali University, People's Republic of China

(Edited by Maryam Tavakol Noorabadi & Kevin D. Hyde)

Published online 5 April 2024