

Outlineoffungi.org – Note 1585 [Fuligo](#)

Web-links: [Index Fungorum](#), [Facesoffungi](#), [MycoBank](#), [GenBank](#)

[Fuligo](#) Haller

According to [García-Martín et al. \(2023\)](#) only four species belong to the clade, which includes the type species of the genus, *F. septica*. Other taxa, described within this genus, including even some varieties of *F. septica* (*F. septica* var. *flava* and var. *candida*), which has been reinstated at species level by [García-Martín et al. \(2023\)](#), are interspersed among other genera of the *Physaraceae*. *Fuligo cinerea* is proposed in the re-erected genus *Aethaliopsis*, *F. laevis* – in the newly proposed genus *Nannengaella* and *F. muscorum* – in the old genus *Lignyidium*.

Reference

[García-Martín J, Zamora JC, Lado C. 2023 – Multigene phylogeny of the order *Physarales* \(*Myxomycetes*, *Amoebozoa*\): shedding light on the dark-spored clade. *Persoonia - Molecular Phylogeny and Evolution of Fungi* 51, 89–124.](#)

Entry by

Anna Ronikier, W. Szafer Institute of Botany, Polish Academy of Sciences, Lubicz 46, 31-512 Kraków, Poland,

Paulina Janik, W. Szafer Institute of Botany, Polish Academy of Sciences, Lubicz 46, 31-512 Kraków, Poland

Published online 23 September 2024