

## Outlineoffungi.org – Note 1587 *Trichamphora*

**Web-links:** [Index Fungorum](#), [Facesoffungi](#), [Mycobank](#), [GenBank](#)

### *Trichamphora* Jungh.

The genus was introduced by Junghuhn (1838) to accommodate *T. pezizoidea*, which forms long-stalked sporophores with discoid-pezizoid sporothecae. The species was later transferred to the genus *Physarum* (Pavillard & Lagarde 1903). In the most recent phylogeny of *Physarales*, *Physarum pezioides* forms a monophyletic clade, therefore García-Martín et al. (2023) proposed a return to the original treatment of the species within the genus *Trichamphora*.

### References

- García-Martín J, Zamora JC, Lado C. 2023 – Multigene phylogeny of the order *Physarales* (*Myxomycetes*, *Amoebozoa*): shedding light on the dark-spored clade. *Persoonia - Molecular Phylogeny and Evolution of Fungi* 51, 89–124.
- Junghuhn FW. 1838 – *Praemissa in floram cryptogamicam Javae insulae*. Batavia, The Netherlands.
- Pavillard J, Lagarde JJ. 1903 – Myxomycètes des environs de Montpellier (1). *Bulletin de la Société Mycologique de France* 19, 81–105.

### Entry by

**Anna Ronikier**, W. Szafer Institute of Botany, Polish Academy of Sciences, Lubicz 46, 31-512 Kraków, Poland,

**Paulina Janik**, W. Szafer Institute of Botany, Polish Academy of Sciences, Lubicz 46, 31-512 Kraków, Poland.

Published online 23 September 2024