

Outlineoffungi.org – Note 1588 [Collaria](#)

Web-links: [Index Fungorum](#), [Facesoffungi](#), [MycoBank](#), [GenBank](#)

[Collaria](#) Nann.-Bremek.

This is an artificial genus with *ca.* five species recognized based on the presence of a collar at the base of the sporotheca ([Lado & Eliasson 2022](#)). However they seem to be unrelated. According to the phylogeny by Fiore-Donno et al. ([2012](#)), the type species of *Collaria*, *C. rubens* (as *Comatricha rubens*) is closely related to *Meriderma* species, while *C. (Comatricha) nigricapillitia* clusters together with *Enethenema*. Also *C. arcyrionema* forms an outermost clade within *Physarales* ([Novozhilov et al. 2022](#)).

References

- [Fiore-Donno AM, Kamono A, Meyer M, Schnittler M et al. 2012 – 18S rDNA Phylogeny of Lamproderma and Allied Genera \(Stemonitales, Myxomycetes, Amoebozoa\). PLoS ONE 7\(4\), e35359,](#)
- [Lado C, Eliasson U. 2022 – Taxonomy and systematics: current knowledge and approaches on the taxonomic treatment of Myxomycetes: updated version. Myxomycetes, 269–324.](#)
- [Novozhilov YK, Prikhodko IS, Fedorova NA, Shchepin ON et al. 2022 – Lamproderma vietnamense: a new species of myxomycetes with reticulate spores from Phia Oắc – Phia Đén National Park \(northern Vietnam\) supported by molecular phylogeny and morphological analysis. Mycoscience 63\(4\), 149–155.](#)

Entry by

Anna Ronikier, W. Szafer Institute of Botany, Polish Academy of Sciences, Lubicz 46, 31-512 Kraków, Poland.

Published online 23 September 2024