

## Outlineoffungi.org - Note 1117 *Chaetophiophoma*

**Web-links:** [Index Fungorum](#), [Facesoffungi](#), [MycoBank](#), [GenBank](#)

### *Chaetophiophoma* Speg.

*Chaetophiophoma* was originally described by Spegazzini (1911), with a single species, *C. trematis*, reported as developing superficially on living leaves of *Trema micrantha* in Argentina. The second species, *C. sorbi*, was described by Li et al. (2020), from *Sorbus* from Italy, as a saprobe on dead branches. An LSU sequence from *C. sorbi* places the genus in *Dermateaceae* (Li et al. 2020). An epitype for *C. trematis* is needed to confirm this taxonomy.

### References

- Li WJ, McKenzie EH, Liu JK, Bhat DJ et al. 2020 – Taxonomy and phylogeny of hyaline-spored coelomycetes. *Fungal Diversity* 100, 279–801. <https://doi.org/10.1007/s13225-020-00440-y>
- Spegazzini C. 1911 – *Mycetes Argentinenses* (Series V). *Annales Mus. Nac. Hist. Nat. Buenos Aires* Ser 3(13), 329–467.

### Entry by

**P.R. Johnston**, Manaaki Whenua – Landcare Research, Private Bag 92170, Auckland 1142, New Zealand

(Edited by **Vinodhini Thiyagaraja, Maryam Tavakol Noorabadi & Subodini N. Wijesinghe**)

Published online 15 May 2023