

Outlineoffungi.org - Note 1161 *Rhynchobrunnera*

Web-links: [Index Fungorum](#), [Facesoffungi](#), [Mycobank](#), [GenBank](#)

Rhynchobrunnera B.A. McDonald, U. Braun & Crous

Rhynchobrunnera was erected by Crous et al. (2021) to accommodate two species of *Rhynchosporium*, *R. lolii* (type species) and *R. orthospora*. A multigene phylogenetic analysis placed *Rhynchobrunnera* sister to *Rhynchosporium*. The two genera are distinguished by their different conidial morphology, *Rhynchobrunnera* possesses 1–3-septate, straight conidia without any apical beak while *Rhynchosporium* species have 1-septate conidia with hooked apical beaks Crous et al. (2021).

Reference

Crous PW, Braun U, McDonald BA, Lennox CL et al. 2021 – Redefining genera of cereal pathogens: *Oculimacula*, *Rhynchosporium* and *Spermospora*. *Fungal Systematics and Evolution* 7(1), 67– 98. <https://doi.org/10.3114/fuse.2021.07.04>

Entry by

Joey B. Tanney, Pacific Forestry Centre, Canadian Forest Service, Natural Resources Canada, 506 Burnside Road, Victoria, BC V8Z 1M5, Canada

(Edited by **Vinodhini Thiyagaraja**, **Maryam Tavakol Noorabadi** & **Subodini N. Wijesinghe**)

Published online 15 May 2023