

## Outlineoffungi.org - Note 1167 *Tricellula*

**Web-links:** [Index Fungorum](#), [Facesoffungi](#), [MycoBank](#), [GenBank](#)

### *Tricellula* Beverw.

Species of *Tricellula*, known only from the asexual state, form Y-shaped, 3-celled conidia. Originally named for a species isolated from air, *T. inaequalis*, the genus has subsequently been found in aquatic habitats ([Van Beverwijk 1954](#), [Webster 1959](#)). ITS sequences from an ex-type culture of *T. inaequalis* (GenBank MH857245, culture CBS 359.53) place the genus in a clade with specimens identified as *Calloria urticae*, the type species of *Calloria* (P.R. Johnston, unpubl. data).

### References

- Van Beverwijk AL. 1954 – Three new fungi: *Helicoon pluriseptatum* n. sp. *Papulaspora pulmonaria* n. sp. and *Tricellula inaequalis* n. gen., n. sp. *Antonie van Leeuwenhoek* 20(1), 1–6. <https://doi.org/10.1007/BF02543704>
- Webster J. 1959 – Experiments with spores of aquatic hyphomycetes: I. Sedimentation, and impaction on smooth surfaces. *Annals of Botany* 23(4), 595–611. <https://doi.org/10.1093/oxfordjournals.aob.a083678>

### Entry by

**P.R. Johnston**, Manaaki Whenua – Landcare Research, Private Bag 92170, Auckland 1142, New Zealand

(Edited by **Vinodhini Thiyagaraja**, **Maryam Tavakol Noorabadi** & **Subodini N. Wijesinghe**)

Published online 15 May 2023