

Outlineoffungi.org - Note 118, [Corylicola](#)

[Corylicola](#) Wijesinghe et al.

[Wijesinghe et al. \(2020\)](#) introduced this genus to accommodate *C. italica* isolated from *Coryllus avellana* L. in Italy. It has similar characters compared to other genera of [Bambusicolaceae](#) [D.Q. Dai & K.D. Hyde](#). These are solitary, scattered, globose to subglobose and ostiolate ascomata; anastomosing and branching pseudoparaphyses; cylindrical asci with a well-developed ocular chamber and short furcate pedicel; and single-septate ascospores. The coelomycetous asexual morph of [Corylicola](#) has holoblastic, phialidic conidiogenous cells and light brown conidia analogous to other members of the family. [Corylicola](#) differs from the other genera of [Bambusicolaceae](#) in having yellowish-brown ascospore masses at the apex of the ascomatal neck (F. Selcuk).

Reference

Wijesinghe SN, Wang Y, Camporesi E, Wanasinghe DN et al. 2020 – A new genus of *Bambusicolaceae* (*Pleosporales*) on *Corylusavellana* (*Fagales*) from Italy. Biodiversity Data Journal 55957. [Doi 10.3897/BDJ.8.e55957](https://doi.org/10.3897/BDJ.8.e55957)