

Outlineoffungi.org - Note 1271 *Infundichalara*

Web-links: [Index Fungorum](#), [Facesoffungi](#), [Mycobank](#), [GenBank](#)

Infundichalara Reblova & W. Gams

Infundichalara is known only from its asexual state. In culture it forms penicillate and *Chalara*-like conidiogenous cells and conidia. DNA sequences from several genes from the ex-type specimen of the type species *I. microchona* (CBS 175.74) place this genus in *Hamatocanthoscyphaceae* (Ekanayaka et al. 2019, Wu & Diao 2023). Koukol (2012) discussed morphological differences between the *Chalara*-like genera *Infundichalara* and *Xenopolyscytalum*.

References

Ekanayaka AH, Hyde KD, Gentekaki E, McKenzie EHC, et al. 2019 – Preliminary classification of *Leotiomycetes*. *Mycosphere* 10, 310–489.

Wu W, Diao Y. 2023 – The *Chalara*-like anamorphs of *Leotiomycetes*. *Fungal Diversity* 119, 213–490.

Koukol O. 2012 – A new species of *Infundichalara* from pine litter. *Mycotaxon* 120, 343–352.

Entry by

P.R. Johnston, Manaaki Whenua – Landcare Research, Private Bag 92170, Auckland 1142, New Zealand

(Edited by **Maryam Tavakol Noorabadi**)

Published online 16 May 2024