

Outlineoffungi.org - Note 1332 *Bezerroplaca*

Web-links: [Index Fungorum](#), [Facesoffungi](#), [Mycobank](#), [GenBank](#)

Bezerroplaca Xavier-Leite, M. Cáceres & Lücking

Bezerroplaca was established under *Gomphillaceae* (*Graphidales*, *Ostropomycetidae*, *Lecanoromycetes*, *Pezizomycotina*, *Ascomycota*) to accommodate five lichenized fungal species based on morphology and phylogeny using SSU and LSU sequence data ([Xavier-Leite et al. 2023](#)). The new genus was typified by *Bezerroplaca lucernifera* (Kalb & Vězda) Xavier-Leite, M. Cáceres & Lücking. In this genus, the thallus is continuous and verrucose with scattered white setae. The apothecia are adnate, emarginate (*Echinoplaca* type), and brown-black. The ascospores are muriform. The hyphophores are setiform and white with a darkened apex. The diahyphae are filiform with spermatozoid terminal segments ([Xavier-Leite et al. 2023](#)).

Reference

Xavier-Leite AB, Goto BT, Lücking R, da Silva Cáceres ME. 2023 – New genera in the lichenized family *Gomphillaceae* (*Ascomycota: Graphidales*) focusing on neotropical taxa. *Mycological Progress* 22(12), 88.

Entry by

Maryam Tavakol Noorabadi, Innovative Institute for Plant Health, Zhongkai University of Agriculture and Engineering, Guangzhou 510225, People's Republic of China

(Edited by **Subodini N. Wijesinghe**)

Published online 31 May 2024