

Outlineoffungi.org - Note 562 [Achrochaeta](#) - morph notes

Web-links: [Index Fungorum](#), [Facesoffungi](#), [MycoBank](#)

[Achrochaeta](#) Réblová & Hern.-Restr. (H)

[Achrochaeta](#) species are saprobes on decaying wood and bark ([Hughes & Kendrick 1968](#), [Réblová et al. 2019](#)). The monotypic genus [Achrochaeta](#) was introduced by [Réblová et al. \(2019\)](#) to accommodate *Chaetosphaeria talbotii*. [Achrochaeta](#) produce hyphomycetous asexual morphs which lack setae and possess cylindrical-clavate conidia ([Réblová et al. 2019](#)). No sexual morph has been reported.

References

- Hughes SJ, Kendrick WB (1968) New Zealand fungi 12. *Menispora*, *Codinaea*, *Menisporopsis*. New Zealand Journal of Botany, 6:323–375. doi.org/10.1080/0028825X.1968.10428818
- Réblová M, Nekvindová J, Kolařík M, Hernández-Restrepo M (2021) Delimitation and phylogeny of *Dictyochoaeta*, and introduction of *Achrochaeta* and *Tubulicolla*, genera nova. Mycologia, 113(2):390–433. doi.org/10.1080/00275514.2020.1822095

Entry by

Rekhani Hansika Perera, Center of Excellence in Fungal Research, Mae Fah Luang University, Muang, Chiang Rai, Thailand.

(Edited by **Kevin D Hyde**)

Published online 26 September 2022