

Outlineoffungi.org - Note 586 [Stilbochaeta](#)

Web-links: [Index Fungorum](#), [Facesoffungi](#), [MycoBank](#)

[Stilbochaeta](#) Réblová & Hern.-Restr.

[Réblová et al. \(2021\)](#) proposed [Stilbochaeta](#) for eight species formerly classified in *Codinaea* and *Dictyochoaeta* which are presumably saprobes. All species are verified by molecular data. The sexual morph of [Stilbochaeta](#) is characterized by perithecial, globose to subglobose, papillate, dark brown, setose ascomata with ostioles, unitunicate, cylindrical-clavate asci, and fusiform, hyaline, septate ascospores, while asexual morph has macronematous, mononematous, brown, unbranched conidiophores, phialidic conidiogenous cells, and falcate, oblong-falcate, ellipsoidal-fusiform, hyaline, aseptate or septate conidia with a setula at each end ([Réblová et al. 2021](#)). A synopsis with diagnostic features were provided by [Réblová et al. \(2021\)](#).

Reference

Réblová M, Kolařík M, Nekvindová J, Réblová K, Sklenář F, Miller AN, Hernández-Restrepo M (2021). Phylogenetic reassessment, taxonomy, and biogeography of *Codinaea* and similar fungi. *Journal of Fungi* 7: 1097, <https://www.mdpi.com/2309-608X/7/12/1097/htm>

Entry by

Ning-Guo Liu, Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai, Thailand

(Edited by: **Kevin D. Hyde**)

Published online 26 September 2022