

Outlineoffungi.org - Note 597 [Rhamphoriales](#)

Web-links: [Index Fungorum](#), [Facesoffungi](#), [Mycobank](#)

[Rhamphoriales](#) K.D. Hyde & Hongsanan

The order [Rhamphoriales](#) was introduced to the class *Sordariomycetes* by [Hyde et al. \(2021\)](#) to accommodate the family *Rhamphoriaceae* which was established by [Réblová et al. \(2018\)](#). *Rhamphoriaceae* consists of three genera, *Rhamphoria* (type genus), *Rhamphoriopsis* and *Rhodoveronaea*. [Rhamphoriales](#) form a distinct clade separate from *Sporidesmiaceae* and are in the range of an order based on divergent time estimation ([Hyde et al. 2021](#)). Species of [Rhamphoriales](#) are saprobes of decaying wood and are reported mainly from temperate regions of Europe ([Hyde et al. 2021](#)). Members of [Rhamphoriales](#) are characterised by ascomycetous sexual morphs and hyphomycetous asexual morphs.

References

- Hyde KD, Bao DF, Hongsanan S, Chethana KWT et al. 2021 – Evolution of freshwater *Diaporthomycetidae* (*Sordariomycetes*) provides evidence for five new orders and six new families. *Fungal Diversity* 107, 71–105. <https://doi.org/10.1007/s13225-021-00469-7>
- Réblová M, Miller A, Réblová K, Štěpánek V 2018 – Phylogenetic classification and generic delineation of *Calyptosphaeria* gen. nov., *Lentomitella*, *Spadicoides* and *Torrentispora* (*Sordariomycetes*). *Studies in Mycology* 89, 1–62 <https://doi.org/10.1016/j.simyco.2017.11.004>

Entry by

Sajeewa S.N. Maharachchimbura, School of Life Science and Technology, Center for Informational Biology, University of Electronic Science and Technology, Chengdu 611731, P.R. China

(Edited by **Kevin D Hyde**)

Published online 26 September 2022