

Outlineoffungi.org - Note 694 *Stromatoneolamyia*

Web-links: [Index Fungorum](#), [Facesoffungi](#), [MycoBank](#)

Stromatoneolamyia Zhurb.

[Zhurbenko \(2021\)](#) introduced this genus to accommodate a sexually typified, lichenicolous taxon, which was reported on *Canoparmelia nairobiensis* from Uganda. The genus is typified by *Z. ugandensis* Zhurb. and it mainly differs from *Neolamyia peltigerae* in 'its well-developed clypeus, completely immersed ascomata clustered in stromatic warts on the host thallus, and the absence of external periphyses'. DNA sequences are lacking thus phylogenetic placement is uncertain in *Ascomycota* ([Zhurbenko 2021](#)).

Reference

Zhurbenko MP. 2021 – Studies on lichenicolous fungi in the Uppsala (UPS) collection curated by the late Rolf Santesson. *Herzogia* 34(2), 493-507. <https://doi.org/10.13158/heia.34.2.2021.493>

Entry by

Nalin N. Wijayawardene, Center for Yunnan Plateau Biological Resources Protection and Utilization, College of Biological Resource and Food Engineering, Qujing Normal University, Qujing, Yunnan, China

(Edited by **Kevin D. Hyde**)

Published online 13 December 2022