

Outlineoffungi.org - Note 697 *Wenhsuisporus*

Web-links: [Index Fungorum](#), [Facesoffungi](#), [MycoBank](#)

Wenhsuisporus C.H. Kuo & Goh

The monotypic dematiaceous hyphomycete *Wenhsuisporus* was proposed by [Kuo et al. \(2022\)](#) for the decaying wood samples collected from submerged in a freshwater stream in Taiwan. The type species, *Wenhsuisporus taiwanensis* C.H. Kuo, S.Y. Hsieh & Goh is characterized by producing slimy masses of hyaline, cylindrical, and aseptate phialoconidia from ampulliform or lageniform conidiogenous cells intercalary in the superficial hyphal network and associated with dark, erect, stiff, unbranched subulate setae bulbous at the base ([Kuo et al. 2021](#)). The phylogenetic analysis based on ITS, LSU and *rpb2* showed that the *Wenhsuisporus* was placed in *Glomerellales* close to *Reticulascaceae*. However, it could not assign any of the families in the order. Therefore, [Kuo et al. \(2022\)](#) treated *Wenhsuisporus* in *Glomerellales* genera *insertae sedis*.

References

Kuo CH, Hsieh SY, Goh TK 2022 – *Wenhsuisporus taiwanensis* gen. et sp. nov., a peculiar setose hyphomycete from submerged wood in Taiwan. *Mycological Progress* 21, 409–426. <https://doi.org/10.1007/s11557-021-01748-y>

Entry by

Sajeewa Maharachchimbura, School of Life Science and Technology, University of Electronic Science and Technology of China, People's Republic of China

(Edited by **Kevin D. Hyde**)

Published online 13 December 2022