

Outlineoffungi.org - Note 964 [Neohormodochis](#)

Web-links: [Index Fungorum](#), [Facesoffungi](#), [MycoBank](#), [GenBank](#)

[Neohormodochis](#) D.P. Wei & K.D. Hyde.

[Neohormodochis](#) was introduced by Wei et al. (2022) to accommodate a monotypic genus with [Neohormodochis septispora](#) D. P. Wei & K. D. Hyde species that selected as the type species. [N. septispora](#) was a saprobe with its asexual morph occurring on dead twig from Kunming City, Yunnan Province, China. Molecular analysis based on concatenated sequences of LSU, mtSSU, ITS and *rpb2* regions supported that [Neohormodochis](#) form a sister clade to *Hormodochies*, while it can be separated from the latter in the white-pruinose conidiomata, crystalline conidiomatal wall, hyaline, holoblastic conidiogenous cells and hyaline, ellipsoidal, dominantly 1-septate, slightly curved conidia (Wei et al. 2022). The sexual morph of [Neohormodochis](#) genus is unknown. The taxonomic placement of [Neohormodochis](#) is in *Stictidaceae*, *Ostropales*, [Lecanoromycetes](#) and *Ascomycota*.

Reference

Wei DP, Gentekaki E, Wanasinghe DN, Hyde KD et al. 2022 – *Neohormodochis septispora* gen. et sp. nov. (*Stictidaceae*) from Yunnan Province, China. *Phytotaxa* 573(2):247–261. <http://dx.doi.org/10.11646/PHYTOTAXA.573.2.5>

Entry by

De-Ping Wei, The Engineering Research Center of Southwest Bio-Pharmaceutical Resources, Ministry of Education, Guizhou University, Guiyang, Guizhou Province 550025, China.

(Edited by **Kevin D. Hyde & Maryam Tavakol Noorabadi**)

Published online 2 April 2024